

# Comment on "Raman spectroscopy study of $\text{Na}_x\text{CoO}_2$ and superconducting $\text{Na}_x\text{CoO}_2 \cdot y\text{H}_2\text{O}$ "

P. Lemmens,<sup>1,\*</sup> P. Scheib,<sup>1</sup> Y. Krockenberger,<sup>2,3</sup> L. Alff,<sup>3</sup> F. C. Chou,<sup>4</sup> C. T. Lin,<sup>2</sup> H.-U. Habermeier,<sup>2</sup> and B. Keimer<sup>2</sup>

<sup>1</sup>*Institute for Physics of Condensed Matter, TU Braunschweig, D-38106 Braunschweig, Germany*

<sup>2</sup>*Max-Planck-Institute for Solid State Research, Heisenbergstr. 1, 70569 Stuttgart, Germany*

<sup>3</sup>*Institute of Materials Science, TU Darmstadt, Petersenstr. 23, D-64287 Darmstadt, Germany*

<sup>4</sup>*Center for Materials Science and Engineering, MIT, Cambridge, MA 02139, USA*

(Dated: February 6, 2008)

The effect of surface degradation of the thermoelectric cobaltite on Raman spectra is discussed and compared to experimental results from  $\text{Co}_3\text{O}_4$  single crystals. We conclude that on NaCl flux grown  $\text{Na}_x\text{CoO}_2$  crystals a surface layer of  $\text{Co}_3\text{O}_4$  easily forms that leads to the observation of an intense phonon around  $700\text{ cm}^{-1}$  [Phys. Rev. B **70**, 052502 (2004)]. Raman spectra on freshly cleaved crystals from optical floating zone ovens do not show such effects and have a high frequency phonon cut-off at approximately  $600\text{ cm}^{-1}$  [Phys. Rev. Lett **96**, 167204 (2006)]. We discuss the relation of structural dimensionality, electronic correlations and the high frequency phonon cut-off of the thermoelectric cobaltite.

PACS numbers: 72.80.Ga, 75.30.-m, 71.30.+h, 78.30.-j

Raman scattering is a well established probe for structural and electronic properties of solids as, e.g. compositional and symmetry information can be gained from the number and frequency of the observed phonon modes [1]. On the other side its high surface sensitivity may also lead to challenges in sample preparation. The cobaltite  $\text{Na}_x\text{CoO}_2 \cdot y\text{H}_2\text{O}$  is a correlated electron system with an enormous thermopower for large  $x > 0.7$  and superconductivity for smaller  $x = 1/3$  and hydration,  $y = 1.3$ . Due to the large mobility of Na on different sites and the mixed nominal Co valency  $\text{Na}_x\text{CoO}_2$  has a complex defect chemistry. In the presence of  $\text{CO}_2$  and humidity surface layers are formed that consists of, e.g.  $\text{CoCO}_3$ ,  $\text{Na}_2\text{CO}_3$  and  $\text{Co}_3\text{O}_4$ . The latter compound is also used as an ingot material in sample preparation [2].

The preparation of large single crystals has been reported from optical traveling floating zone (TFZ) ovens [3] and from NaCl flux ( $\text{NaCl}$ ,  $\text{Na}_2\text{CO}_3$ , and  $\text{B}_2\text{O}_3$  in varying compositions) [4]. TFZ grown crystals can easily be cleaved, while samples from NaCl flux are washed-

out from the flux in water. The latter step may lead to a Na nonstoichiometry. Evidence for degradation and crystallographic changes of  $\text{Na}_x\text{CoO}_2$  and  $\text{Na}_x\text{CoO}_2 \cdot y\text{H}_2\text{O}$  on time scales from minutes to weeks exist in literature. [5, 6].

In a recent Raman scattering investigation of NaCl-flux grown  $\text{Na}_x\text{CoO}_2$  crystals, Shi *et al.* have reported Raman spectra that show 5 phonons with in-plane polarization [7], see Fig. 1, curve a). These modes are attributed to five Raman active modes corresponding to displacements of sodium and oxygen [5]. In contrast to these data Raman scattering investigations on freshly cleaved TFZ grown crystals give only two modes with larger intensity [8]. These modes are attributed to oxygen in-plane and out-of-plane displacements. While the non-observance of the low frequency Na modes is attributed to disorder on the partially occupied Na sites [9], the vibrations of oxygen within the  $\text{CoO}_2$  layers should have characteristic frequencies. Indeed a linear frequency shift of the highest frequency, out-of-plane mode at  $590\text{ cm}^{-1}$  by 5% has been found with increasing Na content in the TFZ crystals [8]. The shift implies that the oxygen modes only weakly depend on the stacking of the  $\text{CoO}_2$  layers and the occupation of Na sites that characterize the ( $\alpha$ ,  $\beta$ ,  $\gamma$  type) crystal structure [9]. With this respect the compound can be considered as two-dimensional and the evolution of electronic correlations with doping dominates the phonon frequency [10]. Our experiments are further supported by recent inelastic X-ray scattering that show a bend over of the highest phonon branch at about  $70\text{ meV} \equiv 583\text{ cm}^{-1}$  [11].

In contrast, the three-dimensional  $\text{Co}_3\text{O}_4$  has a very intense Raman mode at a higher frequency ( $690\text{ cm}^{-1}$ ), i.e. in the same frequency regime as Raman data [7] of NaCl-flux grown  $\text{Na}_x\text{CoO}_2$  crystals. In Fig. 1 we show respective spectra. The small frequency shift and




Figure 1: Raman scattering spectra of (a) NaCl flux grown  $\text{Na}_{0.75}\text{CoO}_2$  at RT (Ref. 7), (b)  $\text{Co}_3\text{O}_4$  at  $T = 200\text{ K}$  and (c) TFZ grown  $\text{Na}_{0.83}\text{CoO}_2$  at  $T = 90\text{ K}$  (Ref. 8).

broadening of curve a) compared to b) is attributed to an oxygen deficiency or a small thickness of the surface layer. Similar data on  $\text{Co}_3\text{O}_4$  have been reported earlier by Hadjiev *et al.* [12] and more recently by Qu *et al.* discussing phase separation [13]. We conclude that the Raman data [7] of NaCl-flux grown  $\text{Na}_x\text{CoO}_2$  are most probably interfered by a degradation of the sample leading to a surface layer of  $\text{Co}_3\text{O}_4$ . We highlight that although from symmetry analysis the same number of Raman active modes are expected, the frequency of the modes in  $\text{Co}_3\text{O}_4$  and  $\text{Na}_x\text{CoO}_2$  differ considerably. The intense  $\text{Co}_3\text{O}_4$  mode at  $690\text{cm}^{-1}$  can be used as a quality measure of cobaltates in thermoelectric applications.

**Acknowledgement:** We acknowledge support by the DFG within the project Le 967/4-1 and the ESF program *Highly Frustrated Magnetism*.

---

\* Electronic address: p.lemmens@tu-bs.de

- [1] See, e.g. contributions to *Light Scattering in Solids*, edited by G. Güntherodt and M. Cardona (Springer, Berlin, 1984-2004).
- [2] At high temperatures ( $T > 850^\circ\text{C}$ )  $\text{Na}_2\text{O}$  has a large volatility. As a result the more stable  $\text{Co}_3\text{O}_4$  forms instead of  $\text{Na}_x\text{CoO}_2$  with smaller or varying  $x$ . At ambient conditions the remaining  $\text{Na}_2\text{O}$  together with  $\text{CO}_2$  and  $\text{H}_2\text{O}$  forms  $\text{Na}_2\text{CO}_3$ .
- [3] F. C. Chou, E. T. Abel, J. H. Cho, Y. S. Lee, J. Phys. Chem. Sol. **66**, 155 (2005); F. C. Chou, J. H. Cho, P. A. Lee, E. T. Abel, K. Matan, and Y. S. Lee, Phys. Rev. Lett. **92**, 157004 (2004); D. P. Chen, H. C. Chen, A. Maljuk, A. Kulakov, H. Zhang, P. Lemmens, and C. T. Lin, Phys. Rev. B **70**, 024506 (2004)
- [4] K. Fujita, T. Mochida, and K. Nakamura, Jpn. J. Appl. Phys. **40**, 4644 (2001); M. Mikami, M. Yoshimura, Y. Mori, T. Sasaki, R. Funahashi and I. Matsubara, Jpn. J. Appl. Phys. Vol. **41**, L777 (2002).
- [5] M.N. Iliev, A.P. Litvinchuk, R.L. Meng, Y.Y. Sun, J. Cmaidalka, C.W. Chu, Physica C **402** (2004) 239-242.
- [6] N. Oeschler, R. A. Fisher, N. E. Phillips, J. E. Gordon, M.-L. Foo, and R. J. Cava, cond-mat/0503690 (2005); M. Karppinen, I. Asako, T. Motohashi, and H. Yamauchi, Phys. Rev. B **71**, 092105 (2005); A. Zorkovský, M. Orendáč, J. Šebek, E. Šantavá, P. Svoboda, I. Bradarić, I. Savić, and A. Feher, Phys. Rev. B **72**, 132412 (2005); C. de Vaulx, M.-H. Julien, C. Berthier, M. Horvatić, P. Bordet, V. Simonet, D. P. Chen, and C. T. Lin, Phys. Rev. Lett. **95**, 186405 (2005); P.W. Barnes, M. Avdeev, J.D. Jorgensen, D.G. Hinks, H.Claus, S. Short, Phys. Rev. B **72**, 134515 (2005); X.N. Zhang, P. Lemmens, B. Keimer, D.P. Chen, C.T. Lin, K.Y. Choi, V. Gnezdilov, F.C. Chou, Physica B **359-361**, 424 (2005).
- [7] Y. G. Shi, Y. L. Liu, H. X. Yang, C. J. Nie, R. Jin, and J. Q. Li, Phys. Rev. B **70**, 052502 (2004).
- [8] P. Lemmens, K.Y. Choi, V. Gnezdilov, E.Ya. Sherman, D. P. Chen, C. T. Lin, F. C. Chou, and B. Keimer, Phys. Rev. Lett **96**, 167204 (2006).
- [9] Q. Huang, M. L. Foo, R. A. Pascal, Jr., J. W. Lynn, B. H. Toby, Tao He, H. W. Zandbergen, and R. J. Cava, Phys. Rev. B **70**, 184110 (2004).
- [10] E. Sherman, M. Fischer, P. Lemmens, P.H.M. van Loosdrecht, G. Güntherodt, Europhys. Lett. **48**, 648 (1999).
- [11] J.-P. Rueff, M. Calandra, M. d'Astuto, Ph. Leininger, A. Shukla, A. Bosak, M. Krisch, H. Ishii, Y. Cai, P. Badica, T. Sasaki, K. Yamada, and K. Togano, Phys. Rev. B **74**, 020504(R) (2006).
- [12] V. G. Hadjiev, M. N. Iliev and I. V. Vergilov, J. Phys. C: Solid State Phys. **21** L190 (1988) and references therein.
- [13] J. F. Qu, W. Wang, Y. Chen, G. Li, and X. G. Li, Phys. Rev. B **73**, 092518 (2006).